



**HAL**  
open science

# Spatial correlations and temporal heterogeneity of the slow dynamics of a colloidal fractal gel

Agnes Duri, Luca Cipelletti

► **To cite this version:**

Agnes Duri, Luca Cipelletti. Spatial correlations and temporal heterogeneity of the slow dynamics of a colloidal fractal gel. 5th international discussion meeting on relaxations in complex systems, Jul 2005, Lille, France. hal-02923760

**HAL Id: hal-02923760**

**<https://hal.inrae.fr/hal-02923760>**

Submitted on 27 Aug 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Agnès Duri and Luca Cipelletti**

Laboratoire des Colloïdes, Verres et Nanomatériaux (LCVN), UMR n°5587, Université Montpellier II, 34095 Montpellier, France


## Experimental System

**Polystyrene colloids** ( $d = 20 \text{ nm}$ ) suspended in a buoyancy-matching  $\text{H}_2\text{O}/\text{D}_2\text{O}$  (45/55 vol/vol)  $\Leftrightarrow \phi_{\text{PS}} = 6 \cdot 10^{-4}$   
+  $\text{MgCl}_2 \Leftrightarrow C_{\text{MgCl}_2} = 10 \text{ mM} \Rightarrow$  **Fractal gel**

## Experimental Set Up

### “Multispeckle” Light Scattering experiments (Single Scattering)

- N°1 **Far field** – Several  $q$  ( $4337 \text{ cm}^{-1} < q < 52177 \text{ cm}^{-1}$ ) – **Time Resolved Dynamics** –  $V \Leftrightarrow$  All the scattering volume  $\Leftrightarrow$  All the speckled image
- N°2 **Imaging geometry** – Single  $q$  ( $q = 10^4 \text{ cm}^{-1}$ ) – **Time and Space Resolved Dynamics** –  $V \Leftrightarrow$  Portion of the scattering volume  $\Leftrightarrow$  Portion of the speckled image



### Time (N° 1) and Space (N° 2) Resolved Degree of Correlation :

$$c_l(t, \tau, \vec{r}) = \frac{\langle I_p(t) I_p(t + \tau) \rangle_{p \in V(\vec{r})}}{\langle I_p(t) \rangle_{p \in V(\vec{r})} \langle I_p(t + \tau) \rangle_{p \in V(\vec{r})}} - 1$$

$I_p(t)$ : intensity p-th pixel at time t

### Spatially Resolved Intensity Correlation Function (N°1):


$$g_2(\tau, \vec{r}) - 1 = \langle c_l(t, \tau, \vec{r}) \rangle_t$$

### Spatial Correlation of the Dynamics (N° 2):

$$X_{cl}(\tau, \vec{r}) = \left\langle \frac{[c_l(t, \tau, \vec{r}) - \langle c_l(t, \tau, \vec{r}) \rangle_t] [c_l(t, \tau, \vec{r} + \Delta \vec{r}) - \langle c_l(t, \tau, \vec{r} + \Delta \vec{r}) \rangle_t]}{\sqrt{\sigma_{c_l}^2(t, \tau, \vec{r}) \sigma_{c_l}^2(t, \tau, \vec{r} + \Delta \vec{r})}} \right\rangle_{\vec{r}}$$

## Experimental Results

### Average Dynamics :




Fit by a “compressed” exponential  
 $g_2(q, \tau) - 1 \propto \exp[-(\tau/\tau_f)^{3/2}]$

$\tau_f \propto q^{-1} \Rightarrow$  Drift motion

$\tau_f \rightarrow t \Rightarrow$  Aging behaviour

Large fluctuations of  $\tau_f$   
 $\Downarrow$   
Heterogeneous dynamics

### Time Resolved Dynamics – Fluctuations Study:




$\rightarrow$  Statistics treatment

$\tau_{\text{max}} \propto \tau_f$

$\sigma^2 c_{\text{max}} \propto q$

$\Downarrow$   
Fluctuations max at large length scale

### Time and Space Resolved Dynamics – Local dynamics Study:



## References :

[1] L. Cipelletti, H. Manley, R.C. Ball, D. Weitz, *J. Phys. Rev. Lett.*, 2000, **84**, 10  
 [2] L. Cipelletti, H. Bissig, Trappe V, P. Ballesta, S. Mazoyer, *J. Phys.: Condens. Matt.*, 2003, **15**, S257- S262  
 [3] A. Duri, H. Bissig, V. Trappe, P. Ballesta, L. Cipelletti, *J. Fluctuations and Noise Letters*, 2005, **5**, 1